



INTEGRATIVE MEDICAL CLINIC

of North Carolina

Contraindications of JOOVV Red Light:

Any of the below described contraindications will require you to use discretion for your own wellbeing.

- Do not use the device if you are photosensitive.
- Do not use the device if using topical, oral, or injectable steroids.
- Do not use the device during the healing period post Lasik eye surgery.
- Do not use the device if pregnant or lactating
- Near Infrared (NIR) at 850 nm may aid hyper-pigmentation; discontinue use if this is a concern.
- Post-surgery users should consult a physician prior to use.
- Users who have had laser hair removal could experience hair regrowth.
- Users with cancer or history of cancer should consult their physician prior to use.
- Users with pre-existing health conditions should consult their physician prior to use.
- Users with hyperthyroidism should consult with their physician prior to use, as use could increase symptoms.
- Users with black pigmented tattoos could experience skin blistering. Blistering can occur by inadvertent heating of the iron oxides and/or the metal salts in the tattoo's black pigment. Tattoo locations can be covered prior to your treatment.
- Users with herpes virus could experience activation of dormant virus with use.
- Users taking medications such as Tetracycline, Digoxin, Retin A, and/or other photosensitive drugs, are recommended to consult with your healthcare provider prior to use.
- Users with a history of facial fillers and Botox injections should consult their dermatologist prior to use.
- Users with breast implants should consult their plastic surgeon prior to use.